



TA: Daniel Destraz

Due on 28th March

Exercise 1 *Binding energy*

a) Show that for a potential of the form $U(R) = -\frac{A}{R^m} + \frac{B}{R^n}$ an equilibrium can only be reached if $n > m$.

b) For a pure van der Waals attraction the potential is often written as

$$U(R) = 4\epsilon \left[\left(\frac{\sigma}{R} \right)^{12} - \left(\frac{\sigma}{R} \right)^6 \right]. \quad (1)$$

Calculate the binding energy (cohesive energy) E_B and the equilibrium distance R_0 .

c) Calculate the effect of thermal expansion, $\Delta R_0(T)/R_0$, on a linear chain of atoms with the potential of part b. Assume that the thermal energy $k_B T \ll E_B$ allows motion of the atoms around the equilibrium position. Think about in what boundaries the atoms can move. From this deduce the average position and compare the result with R_0 .

Hint: Use the expansion $1/(1 \pm \epsilon) \approx 1 \mp \epsilon + \epsilon^2 + \dots$ up to the second order and $\sqrt[n]{1 + \epsilon} = 1 + \epsilon/n + \dots$ for $\epsilon \rightarrow 0$.

Exercise 2 *Madelung constant*

Calculate the Madelung constant for an infinitely long, evenly spaced, linear chain of ions with alternating anions and cations of charge $\pm e$.

Exercise 3 *Linear ionic crystal*

Consider a line of $2N$ ions of alternating charge $\pm q$ with a repulsive potential energy A/R^n between nearest neighbours.

a) Show that the expression for the potential energy can be approximated by

$$U(R) = N \left[\frac{2A}{R^n} - \frac{2 \ln 2 q^2}{4\pi\epsilon_0 R} \right]. \quad (2)$$

b) Show that at the equilibrium separation

$$U(R_0) = -\frac{2Nq^2 \ln 2}{4\pi\epsilon_0 R_0} \cdot \left(1 - \frac{1}{n} \right). \quad (3)$$

c) Let the crystal be compressed so that $R_0 \rightarrow R_0(1 - \delta)$. Show that the work done in compressing a unit length of the crystal has the leading term $\frac{1}{2}C\delta^2$, where

$$C = \frac{(n-1)q^2 \ln 2}{4\pi\epsilon_0 R_0}. \quad (4)$$

Note: Use the complete expression for $U(R)$ instead of $U(R_0)$.