

Abstract EFAS/DGA 2007

Voice discrimination by cochlear implant users

Pyschny, V., Weber, J., Walger, M., von Wedel, H., Meister, H.

Jean Uhrmacher Institute for Clinical ENT–Research, University of Cologne and University ENT-Hospital Cologne

The voice of an individual is an important attribute. Vocal characteristics contribute to differences that help to distinguish one speaker from another. The fundamental frequency (F0), which corresponds to perceived voice pitch, and the vocal tract (VT), which can be described in terms of formant frequencies, are two important measures for voice classification.

A very basic separation of different voices is reflected by the recognition of the speaker gender. Normal-hearing (NH) persons are usually able to identify the speaker gender. In contrast, cochlear implant (CI) patients do not typically have access to the rich spectral and temporal information available to NH listeners and might thus have difficulties to make a decision about the voice gender.

In order to examine the underlying cues F0 and the length of VT respectively were manipulated and it was tested which changes can be recognized. Manipulations were performed using 'Praat' (Boersma et al. 1996) according to a paradigm given in Darwin et al. 2003.

The poster describes the methods used and gives preliminary results for voice gender identification and discrimination for both, CI recipients and normal hearing subjects.

Literatur:

Boersma, P., Weenink, D. (1996): Praat, a system for doing Phonetics by Computer, version 4.4.22. Institute of Phonetic Sciences, University of Amsterdam; www.praag.org.

Darwin, C. J., Brungart, D.S., Simpson, B. D. (2003): Effects of fundamental frequency and vocal-tract length changes on attention to one of two simultaneous talkers. J. Acoust. Soc. Am. 114 (5), 2913 – 2922.

Fu, Q.-J., et al (2005): Voice Gender identification by cochlear implant users: The role of spectral and temporal resolution. J. Acoust. Soc. Am. 118 (3), 1711 – 1718.

